


Firm cooking



Available in organic farming



- High foliar late blight resistance
- Very high storability
- Attractive appearance and excellent cooking quality

General characteristics

Maturity	Second early
Skin / flesh colour	Yellow / Yellow
Tuber shape	Long
Plant height	Medium
Plant growth habit	Semi-upright

Cooking qualities

Cooking type	A	
Flesh firmness	Very good	●●●●●
Discolour. after cook.	None	●●●●●
Frying coloration	Dark	●●○○○

Agronomy

Dry matter content	19 to 20 %	
Tuberisation	15 to 18	
Large tubers number	Medium	
Bruising	Moderat. suscept.	●●●○○
Dormancy	Long	●●●●○
Storability	Long term	●●●●●
De-sprouting	Moderat. suscept.	●●●○○
Internal disorders	Slightly suscept.	●●●●●
Washability	Very good	●●●●●

Pest & disease resistances

Foliar late blight	Resistant	●●●●●
Tuber late blight	Moder. suscept.	●●●○○
PCN Ro 1-4	Quite suscept.	●●○○○
PCN Pa 2-3	Quite suscept.	●●○○○
Common scab	Slightly suscept.	●●●●●
Black dot	Slightly suscept.	●●●●○
Y virus	Moder. suscept.	●●●○○
Yntn virus	Slightly suscept.	●●●●●
Leaf roll virus	Slightly suscept.	●●●●○
Early blight	Slightly suscept.	●●●●○