

Firm cooking



Available in organic farming



- High foliar and tuber late blight resistance
- First early, suitable for early fresh market
- Excellent flavour and cooking qualities

General characteristics

Maturity	First early to early
Skin / flesh colour	Yellow / Pale yellow
Tuber shape	Long oval
Plant height	Medium
Plant growth habit	Spreading

Cooking qualities

Cooking type	A	
Flesh firmness	Very good	●●●●●
Discolour. after cooking	None	●●●●●
Frying coloration	Quite dark	●●○○○

Agronomy

Dry matter content	18 %	
Tuberisation	15 to 18	
Large tubers number	Medium	
Bruising	Moderat. suscept.	●●●○○
Dormancy	Rather short	●●○○○
Storability	Medium	●●●○○
De-sprouting	Quite susceptible	●●○○○
Internal disorders	Slightly suscept.	●●●●●
Washability	Good	●●●●○

Pest & disease resistances

Foliar late blight	Resistant	●●●●●
Tuber late blight	Slightly suscept.	●●●●○
PCN Ro 1-4	Quite suscept.	●●○○○
PCN Pa 2-3	Quite suscept.	●●○○○
Common scab	Slightly suscept.	●●●●○
Black dot	Quite suscept.	●●○○○
Y virus	Moder. suscept.	●●●○○
Yntn Virus	Slightly suscept.	●●●●●
Leaf roll virus	Slightly suscept.	●●●●○
Early blight	Slightly suscept.	●●●●○