


Fresh, export



Available in Organic farming



- High resistance to foliar late blight and to Y virus
- High percentage of large tubers et good storability
- Attractive skin and good washability

General characteristics

Maturity	Early
Skin / flesh colour	Yellow / yellow
Tuber shape	Oval to short oval
Plant height	Medium
Plant growth habit	Semi-upright to spreading

Cooking qualities

Cooking type	A-B	
Flesh firmness	Good	●●●●○
Discolour. after cook.	Light	●●●●○
Frying coloration	Quite dark	●●○○○

Agronomy

Dry matter content	20 to 21%	
Tuberisation	12 to 15	
Large tubers number	High	
Bruising	Quite suscept.	●●○○○
Dormancy	Medium	●●●○○
Storability	Long term	●●●●●
De-sprouting	Quite suscept.	●●○○○
Internal disorders	Slightly suscept.	●●●●○
Washability	Very good	●●●●●

Pest & disease resistances

Foliar late blight	Resistant	●●●●●
Tuber late blight	Quite suscept.	●●○○○
PCN Ro 1-4	Quite suscept.	●●○○○
PCN Pa 2-3	Quite suscept.	●●○○○
Common scab	Moder. suscept.	●●●○○
Black dot	Moder. suscept.	●●●○○
Y virus	Resistant	●●●●●
Yntn virus	Slightly suscept.	●●●●●
Leaf roll virus	Slightly suscept.	●●●●○
Early blight	Slightly suscept.	●●●●○